

AD-A039 803

UNIVERSITY COLL CORK (IRELAND) DEPT OF CHEMISTRY
ENERGY MIGRATION INVOLVING IRRADIATED SOLIDS.(U)
FEB 77 J CUNNINGHAM, D J MORRISEY

F/G 7/4

UNCLASSIFIED

EOARD-TR-77-04

AF-AFOSR-2986-76

NL

1 OF 1
AD
A039803



1 OF 1
AD
A039803



February 28, 1977

FINAL SCIENTIFIC REPORT

AD A 039803

ENERGY MIGRATION INVOLVING IRRADIATED SOLIDS

Period Covered

1.V.1976 - 31.X.1976

Professor Joseph Cunningham,
University College Cork,
Ireland.



DDC
MAY 24 1977
C

DDC FILE COPY

This document has been approved for public release and sale: its distribution is unlimited.

This research has been sponsored in part by [REDACTED]
[REDACTED] the EUROPEAN OFFICE OF AEROSPACE
RESEARCH AND DEVELOPMENT under GRANT NO. AFOSR 76-2986.

AD NO.

Correct
Contract number 6 May 78

is AF-AFOSR-2986-76

per Mrs. Bloce 11:15 ml

Qualified requestors may obtain additional copies from the Defense Documentation Center. All others should apply to the clearinghouse for Federal Scientific and Technical Information.

① REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. Report Number ③ ④ EGARD-TR-77-04 ✓	2. Govt Accession No.	3. Recipient's Catalog Number
4. Title (and Subtitle) ⑥ ENERGY MIGRATION INVOLVING IRRADIATED SOLIDS. ①	5. Type of Report & Period Covered ① Final report <i>rept.</i> 1 May - 31 Oct 1976	
7. Author(s) ⑩ Joseph/Cunningham		6. Performing Org. Report Number ⑨
9. Performing Organization Name and Address Department of Chemistry University College Cork, Ireland ✓		8. Contract or Grant Number ⑮ AF - AFOSR-2886-76
11. Controlling Office Name and Address European Office of Aerospace Research and Development/LNC Box 14, FPO New York 09510		10. Program Element, Project, Task Area & Work Unit Numbers 61102F 9767-99-54 ⑬ ⑭
14. Monitoring Agency Name and Address European Office of Aerospace Research and Development/LNC Box 14 FPO New York 09510		12. Report Date ⑪ 28 Feb 1977
16. & 17. Distribution Statement Approved for public release; distribution unlimited.		13. Number of Pages 42
18. Supplementary Notes		
19. Key Words ENERGY MIGRATION PASSIVATION NITROUS OXIDE IRRADIATED SOLIDS ELECTRON TRANSFER SITES METAL OXIDES		
20. Abstract Results are presented which point to surface sites involving O ₂ ions and/or hydroxyls as important locations for transfer of energy via radiationless process between uv-illuminated metal oxides and probe molecules adsorbed on their surfaces. Attempted passivation of the surfaces by treatment with water vapor enhanced rather than decreased, the extent of interaction between irradiated solid and various probe molecules, including nitrous oxide or methyl halides as electron-accepting probe molecules or C ₂ D ₅ OD as an electron donating probe molecule.		

TECHNICAL SUMMARY

The research carried out under this short low-cost grant mainly represents an extension of a research effort previously funded under AFOSR-71-2148E. Since the Interim and Final Scientific Reports of that preceding grant provide a detailed account of the extensive publications and scientific communications arising from earlier phases of this research, such information is not repeated here, but the earlier reports should be referred to for an overall view of this research effort. In this extension of that research effort experiments have been designed and carried out to examine the exact nature of sites on the surface of irradiated solids which facilitate the transfer of radiation energy to gaseous 'probe' molecules present over the irradiated solids. In line with the agreed statement of work, these studies have included investigations of zinc oxide, magnesium oxide and titanium oxide as irradiated solid substrates; whilst nitrous oxide and alcohols have been used as gaseous probe molecules. This report includes a preprint of a paper describing results of such studies and entitled:


"Reactions Involving Electron Transfer at Semiconducting Surfaces, Part VII. Interactions of Nitrous Oxide and C₂-C₄ alcohols on ZnO and TiO₂ at Room Temperature".

That paper has been submitted for publication and it details evidence to support the conclusion that, for dehydroxylated ZnO or TiO₂ surfaces, the formation of O⁻-type surface intermediates and their subsequent reactions with adsorbed probe molecules (such as aliphatic alcohols) represent a sequence of radiationless processes which effectively convert energy absorbed from incident UV radiation into chemical changes in the adsorbed probe molecules. The preprint also demonstrates that attempted "passivation" of the ZnO and TiO₂ surfaces by treatment with water vapour at 623 K did not affect the reactivity of the dark or uv-illuminated TiO₂ surfaces but did markedly increase the activity of ZnO surfaces for dissociation of nitrous oxide both in the dark and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
214

and additionally under UV-illumination. Although not reported in the preprint, it has also been found that such attempted passivation of MgO surfaces increased their photocatalytic activity for N_2O dissociation, but did so only for photons with $\lambda > 254$ nm. These results upon attempted passivation of ZnO, TiO_2 and MgO surfaces strongly suggest: (i) that surface locations involving a "dangling" surface hydroxyl (which are not known to form on TiO_2 in our conditions) can act as effective sites for transfer of electrons from ZnO or MgO; (ii) that for semiconducting substrates such as Zinc oxides with significant densities (10^{14} - 10^{18} cm^{-3}) of electrons already in the conduction band at room temperature in the absence of uv-illumination, these sites involving dangling hydroxyls allow electron-transfer to probe molecules in the dark; (iii) for insulating substrates such as MgO which lack conduction band electrons, electron transfer becomes possible from sites involving dangling hydroxyls wherever they interact with surface excitons.

The validity of these ideas has been further tested by experiments with methyl halides CH_3Cl and CD_3I , which were selected because, like N_2O , they were reported to be reactive towards electrons but not towards the radical species OH or H. A total absence of photocatalysed dissociation of the methyl halides was noted over uv-illuminated TiO_2 surfaces, which again appeared consistent with absence of hydroxyl-containing surface sites for electron transfer on TiO_2 . On dehydroxylated MgO surfaces photocatalysed dissociation of the methyl halides occurred to methane (80%) plus ethane (20%) but the same surface, after rehydroxylation produced only methane product. The results were consistent with the idea detailed above that presence of surface hydroxyls facilitate reaction of surface excitons with the electron-trapping probe molecules. An important role for surface sites involving hydroxyls has thus been established in energy transfer from irradiated ZnO and MgO to N_2O and methyl halide probe molecules.


J. Cunningham
Principal Investigator

"Reactions Involving Electron Transfer at Semiconducting Surfaces¹:
VII, Interactions of nitrous oxide and C₂ -C₄ alcohols on ZnO and TiO₂
at Room Temperature"


Joseph Cunningham and D.J. Morrissey

Department of Chemistry, University College, Cork. Ireland.


ABSTRACT

Trace amounts of dehydration product desorbed from ZnO or TiO₂ surfaces upon exposure to t-butanol at room temperature whereas alkene products from dehydration of ethanol or isopropanol were only detected following thermally assisted desorption. Dehydration product was not enhanced by nitrous oxide which did, however, increase extent of alcohol dehydrogenation and yield additional nitrogen product from N₂O dissociation. Extent of dehydrogenation and of additional nitrogen formation correlated well with reported reactivity of alcohols towards O⁻ fragments, which are here attributed to N₂O dissociation. Illuminating the Alcohol/Metal Oxide interfaces with photons at 254 nm in a dynamic system with on-line mass spectrometric analysis showed that, with N₂O present, illumination further enhanced dehydrogenation and nitrogen formation. Results are consistent with an important role for coordinatively unsaturated oxygen ion sites in dehydrogenation and with their conversion to surface O⁻ species by hole trapping. Dehydration was largely unaffected by uv-illumination whilst presence of N₂O influenced it in opposite sense to dehydrogenation. Sites active in dehydration are tentatively identified as Lewis acid type metal ion sites on the surfaces.

Earlier papers in this series considered electron-transfer from semiconducting ZnO or TiO₂ to adsorbed molecules, both in terms of *indirect* electron transfer based on collective-electron Band Theory models, and in terms of *direct* electron transfer from isolated electron-donating sites on the ZnO or TiO₂ surfaces^{2,5}. Metal-ion sites possessing lower-than-usual charge and higher-than-usual degree of coordinative unsaturation i.e. $M_{(cus)}^{(n-1)+}$, were identified as sites effective in direct electron-transfer to adsorbed N₂O or O₂. Examples were Ti³⁺ ions adjacent to oxygen vacancies on the surface⁵ of TiO₂ and interstitial Zn⁺_Δ on the surface^{2,3} of ZnO. It was recognised that if equilibration of electrons between such sites and the conduction band of these n-type semiconducting solids occurred rapidly via. I,




then the electron-donating power of these surface metal-ion sites should vary with position of the Fermi level. This prediction was confirmed^{2,5} by careful measurement of the very small amounts of nitrogen product evolved from surfaces of various zinc oxides by dissociative-electron-attachment to nitrous oxide at room temperature according to 2,



Other lines of evidence have also indicated electron exchange between bulk and surface locations of zinc oxides^{6,7} but evidence has been lacking for corresponding exchange processes in TiO₂. Surfaces of TiO₂ powders have, however, been shown to donate^{8,9,10} or accept^{11,12} electrons from adsorbed organic molecules in certain conditions.

The present work extends to aliphatic alcohols our approach of utilising room-temperature reaction of surfaces with appropriate probe

molecules as a means of assessing number and nature of highly active surface sites. This extension of the approach was prompted by, and was designed to test, the following ideas advanced by previous workers in relation to surface activity for dehydrogenation and dehydration of aliphatic alcohols: *firstly*, the idea that primary and secondary alcohols are good reagents for electronic holes at surfaces of ZnO, either through direct interaction^{1,3,14} or after hole capture by OH^{15,16}. The validity of these suggestions was of particular interest since, if correct, they would establish that alcohols could be used as hole-probes in similar manner to that already demonstrated for N₂O as an electron-probe in examining charge-localization at these semiconductor surfaces. Since concentrations of surface-localised holes were expected to be low for our n-type ZnO and TiO₂ samples in the dark at room temperature, present experimental tests of these suggestions include an examination of the interactions of alcohols with uv-illuminated samples having large concentrations of photogenerated holes. Such electronic holes in an array of lattice O²⁻ ions should possess some of the characteristics of O⁻ species⁷; *Secondly*, the idea that O⁻ exhibits selective reactivity towards primary and secondary, but not tertiary, alcohols, as previously postulated by Warman^{1,7,18}. Warman further suggested that, when N₂O is used as the source of O⁻ species, secondary reactions 3a and 3b occur in the presence of primary or secondary alcohols and produce additional dissociation of N₂O.



Insufficient information prevented identification of the ionic intermediates, (X+Y)⁻, in the gas-phase reactions studied by Warman. However,

it was hoped that examination of analagous reactions on ZnO and TiO₂ surfaces in the present study would reveal surface processes similar to 3a and 3b and might clarify the nature of ionic intermediates and products on the metal oxide surfaces.

EXPERIMENTAL

Materials: Powdered, high-purity metal oxides were identical to the ZnO (SP500) and TiO₂ (Rutile MR-128) used in previous studies²⁻⁵ and were obtained by courtesy of New Jersey Zinc Co. Thin polycrystalline layers on quartz substrates were prepared and activated at 623K as previously described. Alcohols were Analar reagent grade dried over freshly dehydrated molecular sieve and purified by trap-to-trap distillations, except for anhydrous C₂D₅OD or "C₂D₅OD + 5% D₂O" which were used as supplied by Stohler Isotopes. High purity nitrous oxide, BOC Grade X, was used as obtained.

Procedures for non-illuminated samples: Observations on the gas phase above non-illuminated samples were made with the closed quartz reactor and associated greaseless high-vacuum system depicted in Fig.1a, wherein products were detected from room temperature interactions of alcohols and/or nitrous oxide with metal oxides previously activated for 16 hr. in vacuo at 623K. Products non-condensable (n.c.) at 77K were measured with Pirani or Ionization gauges and identified with a CEC type 21-620A mass spectrometer during periodic freezing-out of reactants into a trap at 77K. Since such measurements could only monitor growth of those products which desorbed from the catalyst surface at room temperature, a sequence of "desorption/pressure - measurement/mass analysis" was carried out with the system shown in Fig.1a while the catalyst, which had been exposed to

Alcohol and/or nitrous oxide at room temperature, was heated in vacuo by 50° increments to 623K. Electron spin resonance spectra were taken with a Decca X-1 spectrometer or a Varian model-, upon TiO_2 samples to which alcohol was admitted as a vapour at room temperature.

Photoassisted Surface Reactions: Interfaces between Metal Oxide sample and Alcohol or (Alcohol+N₂O), which had completed the fast 'dark-reaction' characteristic of initial 0.5h contact times⁵, were then illuminated by the low intensity output of a 15-watt low pressure Hg-arc lamp with emission mainly at 254 nm. Lamp configuration relative to the sample within an MgO-coated cavity was arranged to limit any possible surface heating effect to <5° temperature rise which, if induced thermally, did not significantly disturb the dark-equilibrated behaviour. Growth of any product which desorbed into the gas-phase at room temperature was followed and analysed over periods of illumination up to 6 hr. Photon flux at the sample position was determined by calibration with potassium ferrioxalate actinometer. Results obtained at illuminated interfaces are indicated by use of an asterisk e.g. N₂O/ ZnO^* or C₂H₅OH/ TiO_2^* . Photoassisted products were monitored *either* in the same system as for dark reaction (see Fig. 1a) *or* more directly in the dynamic flow system depicted in Fig. 1b. In this latter system, gases emerging from the reactor at pressures ca. 10^{-4} N m⁻² entered into the ion source of a Micromass 6 mass analyser located 15 cm from the catalyst sample via. a high conductance path. Photoassisted changes in composition of the gas flow could thus be very rapidly detected and monitored.

RESULTS AND DISCUSSION

Section I: Formation and Desorption of Products in static reactors.

I.1. Nitrogen product from N_2O/ZnO and $(N_2O + Alcohol)/ZnO$ interfaces at room temperature:

Figure 2A(i) illustrates the rapid growth of gaseous product non-condensable at 77K which occurred during the first hour of contact at room temperature between freshly activated ZnO and nitrous oxide. Plot (i) of Fig. 2B demonstrates that formation of noncondensable product over the same ZnO surface in the dark at room temperature was slightly reduced when t-butanol was premixed with N_2O prior to contacting the freshly activated surface. Plots (ii) of Fig. 2A and 2B demonstrate, however, that the limiting yield of noncondensable product, $V(N_2)_{lim}$ as observed at room-temperature contact times >60 min, was greatly enhanced when either ethanol or isopropanol was premixed with N_2O in 1:1 mole ratio and simultaneously contacted with the metal oxide. Mass spectrometric analysis of the noncondensable product established that it was $>98\%$ nitrogen and that no nitric oxide or oxygen was detectable. Empirical kinetic analysis showed that the increase in nitrogen product, $V(N_2)_t$ detected after dark contact time t , obeyed the first-order type expression,

$$\log \left(\frac{V(N_2)_{lim}}{V(N_2)_{lim} - V(N_2)_t} \right) = k_d t \quad (4),$$

for contact times 5 - 30 mins. Table I summarises the values of $V(N_2)_{lim}$ and k_d which gave the "best fit" of data to this expression. These data point to two significant differences between the influence of tertiary as distinct from primary or secondary alcohol at the N_2O/ZnO interface:

(a) Nett nitrogen product formation from nitrous oxide at contact times

TABLE I

Limiting yields of Nitrogen Produced and desorbed over ZnO at room temperature

SYSTEM	$P_0(P_{\text{React}})$ N_m	$V(N_2)_{\text{Lim}}$ $\text{mls/m}^2 \times 10^{-5}$	$t_{\frac{1}{2}}$ (Min)	θ_0	k_d (Min^{-1})
N_2O/ZnO	45.0	4.16	8.1	0.011	8.6×10^{-2}
$N_2O + C_2H_5OH/ZnO$	45.0 (each)	7.73	7.7	0.019	9.0×10^{-2}
$N_2O + (CH_3)_2CHOH/ZnO$	45.0 (each)	7.76	8.4	0.019	8.3×10^{-2}
$N_2O + (CH_3)_3COH/ZnO$	45.0 (each)	3.53	4.1	0.009	16.9×10^{-2}

>1 hr., $V(N_2)_{lim}$, was approximately doubled by premixing ethanol or isopropanol with the nitrous oxide admitted to the freshly activated ZnO surface; whereas admixture of t-butanol slightly reduced $V(N_2)_{lim}$; (b) the apparently first-order rate constant, k_d , for dissociation of N_2O to N_2 at the 'dark' ZnO surfaces was not significantly affected by premixed ethanol or isopropanol, whereas t-butanol increased k_d , despite its overall inhibiting effect on $V(N_2)_{lim}$.

The effects noted for ethanol and isopropanol so strongly resemble those reported by Warman^{1,2,10} (see Introduction) that a similar interpretation was suggested, viz. that initial dissociation of N_2O occurred via (2) on the metal oxide and that the resultant $O^- (ads)$ species reacted rapidly with alcohol, producing surface intermediates, $(X+Y)^-$, which in turn rapidly dissociated other N_2O molecules via 3a and 3b. Such a mechanism, with process 2 containing the slow rate-determining step, would account for the overall rate constant k_d remaining unaffected by ethanol or isopropanol whilst the nitrogen yield $V(N_2)_{lim}$ doubled, since each type (2) event leads eventually via 3 to dissociation of at least two N_2O molecules. Our observation that t-butanol did not increase $V(N_2)_{lim}$ appeared consistent with Warman's conclusion that tertiary alcohols do not undergo type 3 reaction with O^- , and hence cannot lead to additional N_2O dissociation.

Data points to the right of an arrow on each plot in Fig. 2 show that illumination by UV light mainly at 254 nm caused additional photo-assisted dissociation to nitrogen product at ZnO interfaces which previously had attained their limiting value of $V(N_2)_{lim}$ in the dark. Two

contributions to this photoassisted dissociation can be distinguished on the traces in Fig. 2A: a slow linear growth of $V(N_2)^*$ with duration of illumination, which was the sole photoassisted process for N_2O/ZnO^* ; and an initial rapid photoassisted growth $V(N_2)^*$, of nitrogen persisting only during the initial $\frac{1}{2}$ hr. of illumination at the $(N_2O + EtOH)/ZnO^*$ interface. For this latter interface the rate of photoassisted N_2 formation returned to that characteristic of N_2O/ZnO^* interfaces at illumination times $>\frac{1}{2}$ hr.

Data to the right of the arrows in Fig. 2B for mixtures of N_2O with isopropanol or t-butanol over the illuminated ZnO surface, also appear consistent with contributions by both an initial fast photoassisted process and a slower continuing process to growth of $V(N_2)^*$. Mass spectrometric analysis demonstrated that the apparent increase in n.c. product from illuminated t-butanol/ ZnO^* interface shown in Fig. 2B originated from photoassisted formation of methane and not N_2 . This was the only illuminated interface found to yield appreciable desorption of methane product at room temperature and the significance of this observation is discussed in a later section.

Previous infra-red studies indicated a high degree of dehydroxylation for ZnO surfaces pretreated in vacuo at the 623-673K temperatures employed in the present study.¹⁹ Isotopic composition of the methanes produced at such ZnO surfaces by interaction with CDI was likewise previously shown to be consistent with extensive dehydroxylation of ZnO activated at these temperatures.⁴ Consequently the data in Fig. 2 for N_2 formation from N_2O/ZnO interfaces approximate to the situation for a dehydroxylated surface. It

TABLE 2

Limiting Yields of Nitrogen Produced and Desorbed over TiO_2 at Room Temperature

SYSTEM	P_{O_2} (React) Nm^{-2}	$V(\text{N}_2)_{1\text{atm}}$ $\text{mls}/\text{m}^2 \times 10^{-4}$	$t_{\frac{1}{2}}$ (Min)	θ_{O}	k_d (min^{-1})
$\text{N}_2\text{O}/\text{TiO}_2$	45.0	2.34	11.5	0.057	6.02×10^{-2}
$\text{N}_2\text{O} + \text{C}_2\text{H}_5\text{OH}/\text{TiO}_2$	45.0 (each)	6.90	5.1	0.176	1.36×10^{-1}
$\text{N}_2\text{O} + (\text{CH}_3)_2\text{CHOH}/\text{TiO}_2$	45.0 (each)	6.89	4.7	0.174	1.49×10^{-1}
$\text{N}_2\text{O} + (\text{CH}_3)_3\text{COH}/\text{TiO}_2$	45.0 (each)	2.01	5.7	0.049	1.23×10^{-1}

was of interest to rehydroxylate such ZnO surfaces and again observe extent of nitrogen formation, in view of suggestions that surface hydroxyls on metal oxides represent active surface sites. The ZnO surfaces were rehydroxylated by baking in water vapour at 623 for 2 hrs. after which water was pumped away at 623 and the sample cooled to room temperature. The growth of nitrogen product over this rehydroxylated surface in the absence of illumination is represented by curve(ii) of Fig.3A. Comparison with the initial 80 min. of curve(i), which was obtained on the same ZnO layer prior to its rehydroxylation, leaves no doubt that rehydroxylation yielded additional surface sites on ZnO which were active for N_2O dissociation to N_2 at room temperature in the static reactor. Curve(iii) of Fig.3A illustrates that UV illumination of the interface between N_2O and rehydroxylated ZnO caused a further large enhancement in the rate of N_2O dissociation. Thus it is clear that rehydroxylated ZnO surfaces were much more active than dehydroxylated ZnO for N_2O dissociation both in the dark and particularly under UV illumination.

1.2. *Nitrogen product from N_2O/TiO_2 and $(N_2O+Alcohol)TiO_2$ interfaces at room temperature;*

Data to the left of the arrows in Figure 4 illustrates growth of nitrogen product during the first hour of contact of freshly-activated non-illuminated TiO_2 with N_2O or with $(N_2O+Alcohol)$. Empirical kinetic analysis again established that data for contact times 5-30 min. could be adequately fitted to the first-order type expression 4. Table 2 lists values so determined for $V(N_2)_{lim}$ and k_d over the TiO_2 surface. Inspection of the $V(N_2)_{lim}$ values reveals that, at room temperature, t -butanol failed to enhance nitrogen product formation at the dark N_2O/TiO_2 interface, whereas admixture of ethanol or isopropanol greatly enhanced $V(N_2)_{lim}$ relative to that with only N_2O present. Comparison

of data in Table 2 with Table 1 reveals that admixtures of ethanol or isopropanol with N_2O trebled the limiting yield of nitrogen product $V(N_2)_{lim}$ observed when contacted with TiO_2 , in contrast to the doubling upon contact with ZnO . Since enhancement by a factor of 3 was just that noted by Warman^{17, 18} for reaction with gas-phase O^- , this observation suggested that O^- intermediates on TiO_2 surfaces reacted with these alcohols with stoichiometry similar to that reported for processes 3a and 3b in the gas phase. The smaller enhancement of (N_2) noted in Table I for corresponding ZnO interfaces could arise *either* from loss of some fraction of surface O^- radicals into the ZnO bulk by hole migration (which recent results of Lundsford support⁷) or by the blocking of some active sites on ZnO through rapid and irreversible chemisorption of alcohol (see below).

In contrast to ZnO interfaces, at which only t-butanol increased the apparently first-order rate constant k_d for N_2 formation from N_2O , all the alcohols caused k_d over $(N_2O+Alcohol)/TiO_2$ interfaces to increase to similar values approximately double those over N_2O/TiO_2 . Unchanged values for $V(N_2)_{lim}$ with admixtures of t-butanol gave no evidence for occurrence of processes 3a or 3b with this alcohol. Consequently it appeared that only process 2 could contain a slow r.d.s. common to the three systems and that each alcohol enhanced this r.d.s. to comparable extent on TiO_2 . Process 2 involves the sequence:- N_2O adsorption dissociation of N_2O within the surface; and finally, N_2 desorption. Data additional to those on rate of N_2 appearance in the gas phase would be necessary to determine which of these steps was enhanced by the alcohols to give rise to observed increases in k_d . Evidence on

this question is discussed in a later section.

Data to the right of an arrow in Fig. 4 were taken with photons of $\lambda=254$ nm incident onto the previously dark-equilibrated interfaces. The lack of any significant photoassisted dissociation to N_2 at the N_2O/TiO_2^* interface argued against significant photocatalysis of Process 2 at that interface. Consequently no photoassisted formation of additional $O_{(ads)}^-$ intermediates from N_2O was thought to occur at N_2O/TiO_2^* interfaces. Plots 4A(ii) and 4B(ii) demonstrate, however, that readily measurable rates of N_2 production were observed when ethanol or isopropanol were simultaneously present with N_2O at the illuminated TiO_2 surface. A working hypothesis capable of explaining these observations was that illumination of the TiO_2 surface directly produced surface $O_{(ads)}^-$ intermediates without the need to photodissociate N_2O , (e.g. by trapping photo-generated holes at pre-existing coordinatively unsaturated O^{2-} ions²⁰). Reaction of adsorbed alcohol with such O^- entities via 3a to produce secondary surface radicals capable of reacting with N_2O via 3b, would account for the additional N_2 observed upon illumination despite the absence of O^- fragments from N_2O photodissociation. Absence of any photoassisted nitrogen product formation in the presence of t-butanol (see plot 4B (i) to the right of the arrow), is consistent with reported low reactivity of O^- towards this tertiary alcohol (recall that the photoassisted increase in noncondensable products from t-butanol/ TiO_2 shown in Fig. 4 B plot (i) was due to formation of methane and not N_2).

Negative results of experiments aimed at establishing whether rehydroxylation of the TiO_2 surfaces enhanced N_2O dissociation to nitrogen are summarised in Fig. 3 B, which shows that growth of N_2 product followed

almost identical curves for dehydroxylated and 'rehydroxylated' TiO_2 surface previously activated in vacuo at 623 for 16 hr. was carried out in the same manner as for ZnO (viz. 2 hrs. in water vapour at 623K) but figure 5.b. makes clear that this treatment caused neither any increase in N_2 formation in the dark nor any photoassisted N_2O dissociation. This result makes it improbable that the enhanced dissociation of N_2O here observed in the presence of alcohol over the non-illuminated TiO_2 surface could have arisen from any rehydroxylation of the surface by the alcohol. However, some workers report lack of success in attempts to rehydroxylate TiO_2 surfaces²¹ and the conclusion that surface hydroxyls on TiO_2 are inactive for N_2O dissociation must remain tentative until direct evidence becomes available on the extent of surface rehydroxylation achieved with H_2O or alcohol. Results of studies on the interaction of N_2O and alcohols over ZnO or TiO_2 surfaces obtained on static systems by other techniques are described in the following paragraphs.

I. 3. *Products released by thermally-assisted desorption from ZnO or TiO_2 interfaces previously exposed to alcohol and/or N_2O .*

The t-butanol/ ZnO interface was the only case in which release of product into the gas phase accompanied room-temperature adsorption of alcohol onto ZnO or TiO_2 surfaces previously activated in vacuo at 623°K (cf. Table 3). Our failure to detect aldehyde or ketone products from the other alcohols in the gas phase above the surfaces of these n-type semiconducting solids did not appear fully consistent with previous claims that a high Fermi level should promote dehydrogenation^{22, 23}.

TABLE 3

Extent of Room-temperature Product formation at Alcohol Vapour/Metal Oxide Interfaces - as deduced from thermal desorption in vacuo.

SYSTEM	Reaction Conditions at 390K	DESORBED PRODUCTS ^a			
		(-H ₂ O)	(-H ₂)	(Diss)	H ₂ (g)
Alcohol/Metal Oxide					
C ₂ H ₅ OH/ZnO ^e	dark	4.7	-	1.4 ^b	17.0
"	+hv	5.0	-	1.9 ^b	18.0
"	+N ₂ O	3.2	-	2.6 ^b	14.0
"	hv+N ₂ O	4.4	-	2.7	15.0
C ₂ H ₅ OH/TiO ₂ ^f	dark	19.0	-	13.0 ^b	0.6
"	+hv	19.0	-	4.1 ^b	2.1
"	+N ₂ O	5.8	-	30.0 ^b	0.2
"	hv+N ₂ O	2.6	-	12.0 ^b	-
(CH ₃) ₂ CHOH/ZnO ^e	dark	3.9	1.9	-	16.0
"	+hv	3.8	2.2	-	11.0
"	+N ₂ O	2.8	2.8	-	8.7
"	hv+N ₂ O	2.5	5.7	-	7.3
(CH ₃) ₂ CHOH/TiO ₂ ^f	dark	69.1	-	6.8 ^c	-
"	+hv	58.0	0.5	14.0	-
"	+N ₂ O	47.0	0.2	4.5 ^c	-
"	hv+N ₂ O	44.0	1.6	9.9	-
(CH ₃) ₃ COH/ZnO ^e	dark	16.0	-	-	CH ₄
"	+hv	18.0	-	0.02 ^d	-
"	+N ₂ O	16.0	-	0.01 ^d	-
"	N ₂ O+h	17.0	-	0.12 ^d	-
"	dark	0.06 ^g	-	-	-
"	+hv	0.09 ^g	-	0.06 ^g	0.07 ^g
"	N ₂ O+hv	-	-	0.20 ^g	0.09 ^g
(CH ₃) ₃ COH/TiO ₂ ^f	dark	81.0	-	- ^d	-
"	+hv	100	-	1.5	-
"	+N ₂ O	70.0	-	-	-
"	hv+N ₂ O	100	-	3.4	-
"	hv	4.2 ^g	-	-	0.10 ^g
"	N ₂ O+hv	1.7 ^g	-	-	0.15 ^g

a) Normalised to μl of product gas (STP) desorbing per m^2 of oxide surface.

b) Detected as butadiene + CO₂ + methyl acetylene + propene.

c) Detected as acetaldehyde.

d) Detected as acetone.

e) Alkene and H₂ desorbed 300-460° but aldehyde on ketone 460-620K.

f) All product except acetone desorbed 300-510K but acetone desorbed 460-620K

g) Desorbed at room temperature.

Efforts were therefore made to detect the expected dehydrogenation products by thermal desorption after gas phase and any physisorbed alcohol had been pumped away at 300K away under continuous evacuation at $< 10^{-4}$ Nm⁻².

Data on these effects, which are summarised in Table 3, do not provide reliable absolute values for surface densities of sites active for alcohol dehydration or dehydrogenation at room temperature, because of possibilities for additional reaction during the thermal desorption procedure. They do, however, provide 'baseline' values for comparison with amounts of product thermally desorbed from the same interfaces in different conditions e.g., when exposed to uv-illumination or to nitrous oxide admixed with the alcohol.

Significant numbers of 'active sites' capable of alcohol dehydration were indicated by thermal desorption of corresponding alkene in readily measurable amount, (tabled under -H₂O in Table 3). This was true for both ZnO and TiO₂, but the specific activity of TiO₂ was greater in all cases. For the TiO₂ surface, extent-of-dehydration increased in the sequence, t-butanol>isopropanol>>ethanol, which is the same as that usually observed for alcohol dehydration by acid-type centres in homogeneous or heterogeneous catalysis. Appearance of the olefine product at temperatures much below those previously reported indicates that some at least, of the dehydration sites were of unusually high activity.

As expected, no dehydrogenated product was obtained for t-butanol but the ratio,(dehydrogenation/dehydration),increased for isopropanol and ethanol. Thermally assisted desorption of the anticipated dehydrogenation products,(acetone or acetaldehyde from isopropanol or ethanol respectively),occurred to lesser extent than release of dehydrated products or of molecular hydrogen at temperatures $<498\text{K}$. The histograms in Fig. 5 illustrate the higher desorption temperatures required for release of acetone product than for propene or hydrogen desorption from the isopropanol/ZnO interface. The acetone histogram also represents (by the fully blackened sections) the thermal desorption behaviour noted for the same surface when acetone rather than alcohol was initially absorbed at room temperature. At the higher temperatures required for desorption of aldehydes or ketones, thermally-assisted rearrangements occurred - as evidenced by partial conversion of preadsorbed acetone to isobutene or of preadsorbed acetaldehyde to 1, 3 butadiene. Extent of ethanol dissociation corresponding to products related to adsorbed acetaldehyde is tabled under (Diss) in Table 3. This latter observation contrasts with a report by McArthur²⁴ that acetaldehyde desorbed readily from ZnO and the difference suggests that ethanol selectively displaced acetaldehyde at the temperatures employed in McArthur's study.

Admixture of nitrous oxide with the ethanol admitted to dark ZnO or TiO₂ caused a definite increase in products attributable to acetaldehyde (which then suffered further degradation

upon thermal desorption). This observation, and a smaller increase in acetone from isopropanol, appeared consistent with the working hypothesis developed in Section I, according to which nitrous oxide dissociated on the surfaces to yield O^- fragments which enhanced dehydrogenation by then abstracting α -hydrogen from primary or secondary alcohol. Data in Table 3 also show that dehydration products experienced a decrease when N_2O was admixed with primary or secondary alcohols, in contrast to the increases which N_2O brought about in dehydrogenation.

A slight increase in extent of dehydration was observed for t-butanol under illumination at either ZnO or TiO_2 surfaces but nitrous oxide did not further enhance this photodehydration. Comparison of the first and second data rows for other Alcohol/Metal Oxide systems in Table 3 reveals that no significant enhancement of dehydrated or dehydrogenated product above the baseline values resulted from illumination of the interfaces through a quartz wall with the output of a 15 watt low pressure mercury arc lamp. Photons of λ 254 nm predominated in this output and such photons had sufficient energy to create holes and electrons in surface and sub-surface regions of the ZnO or TiO_2 layers. Reports by previous workers, to the effect that alcohols reacted with electronic holes at uv-illuminated ZnO surfaces, had led us to expect additional oxidation of alcohols at that illuminated interface relative to the baseline values observed in the dark. Our failure to detect ^{such} addition-

al photoassisted dissociation of ethanol or isopropanol vapour casts serious doubts on the idea that electronic holes promote alcohol oxidation. The last data line for the t-Butanol/Metal Oxide system in Table 3 supports a similar negative conclusion, even when nitrous oxide and t-butanol were simultaneously present over these uv-illuminated interfaces. Particular importance attaches to this latter observation because N_2O represented a reducible, and t-butanol an oxidisable, reagent. In terms of electronic theories²², photogenerated holes and hole-hole-electron pairs should promote their photoreduction and photooxidation respectively. Consequently this negative result argues against an important role of electronic factors at the interfaces between illuminated ZnO or TiO_2 and (t-Butanol + nitrous oxide).

1.4. ESR measurements:

Further evidence of the small influence of adsorbed alcohol upon electron localization/delocalization at the metal oxide interface emerges from the ESR spectra shown in Fig. 6. Spectrum (i) was measured at 77K on a sample of TiO_2 which had been outgassed in vacuo at 623K and it corresponds well to broad anisotropic adsorption band usually attributed to paramagnetic Ti^{3+} centres at the TiO_2 surface. Spectrum (ii) was not significantly altered if the sample was warmed in vacuo to 300K and again cooled to 77K but a new spectrum (ii) resulted if such samples were exposed to the saturation vapour pressure of

ethanol or sec-2-butanol for 10 min. and re-evacuated at 300K before recooling to 77K. When such samples were subsequently heated to 623K under continuous evacuation and remeasured at 77K spectrum (i) was again obtained. The reversible nature of the change brought about by brief exposure to alcohol vapour at room temperature would be consistent with chemisorption of the alcohol. It must however be concluded from the small differences between spectra (i) and (ii), either in respect of integrated intensity or magnetic parameters, that surface paramagnetic Ti^{3+} centres were not markedly affected by the chemisorbed alcohol. The changes in shape of the resonance would be consistent with weak dipolar interaction between chemisorbed alcohol and Ti^{3+} , resulting in a slight increase in g_{11} of the latter.

The absence of any marked change in integrated intensity of the spectrum certainly would not be consistent with extensive charge transfer between alcohol and the vacuum-activated TiO_2 surface. Thus if chemisorbed alcohol acted as a donor increase in Ti^{3+} centres would be expected. Conversely decrease in their number should have occurred if Ti^{3+} centres transferred electrons to chemisorbed alcohol.

SECTION II

Reactions of alcohols and/or (Alcohol+N₂O) mixtures at illuminated ZnO and TiO_2 surfaces under dynamic conditions.

Since results in Section 1 showed that aldehyde and ketone products from alcohol dehydrogenation reactions remained adsorbed on non-illuminated ZnO and TiO_2 surfaces in the static

reaction system and were only gradually desorbed on heating to 625°K in vacuo, experiments were carried out in dynamic conditions to determine if products of photoassisted elimination reactions then remained adsorbed in similar manner, or if they were released to the gas phase at room temperature. Oxide samples were therefore prepared and pre-treated as described above and then exposed to low dynamic pressures of reactant gas(es) at ca 10^{-4} Nm⁻², which were established over the samples as a balance between inlet leak-rate of reactants through a metal variable leak valve and their removal by a continuously operating ion-pump with pumping speed of 50 litres sec⁻¹. A micromass 6 mass analyser located 15 cm from the oxide surfaces along a high conductance path was employed for on-line gas analysis. Reactant alcohols, or (alcohol+N₂O) mixtures in approximately 1:1 mole ratio, were mass-analysed at various time intervals, prior to the commencement of illumination, until reproducible reactant spectra were obtained. Continuous illumination of the GAS/SOLID interface was then commenced and the gaseous phase mass-analysed at various times during an illumination period of approximately three hours. Interfaces were illuminated through a quartz envelope by the unfiltered output of a 15-watt low pressure Hg lamp with output concentrated principally at 254 nm. and results are summarised in Figures 7A and 7B.

II. 1. Photoproducts from Isopropanol/Metal Oxide interfaces:

On-line mass analysis provided no evidence for measurable enhancement of peaks at m/e values corresponding to propene, the dehydration product detected from ZnO interfaces by thermal desorption (cf. Section 1). Exposure of the illuminated ZnO surface

to alcohol was, however, several orders of magnitude less in these on-line analyses (ca 4×10^{-3} torr sec) than for typical exposures (ca 10^3 torr sec) in the static reactor experiments.

The lower curve of Fig. 7A illustrates photo-assisted production of acetone dehydrogenation product upon illumination of an *isopropanol/ZnO* interface through quartz. Note the rapid initial increase in gas-phase acetone, reaching a maximum approximately 4 minutes after the commencement of uv illumination. This increase was followed by a slow decrease in dynamic pressure of acetone until, at times ~ 170 min. after the commencement of illumination, photoproduction of acetone had ceased. These results indicated that, as illumination proceeded in the presence of isopropanol, surface sites capable of photoassisted dehydrogenation were progressively removed. The results would be consistent with photogeneration of surface O^- sites and their reaction with isopropanol to yield dehydrogenated product plus inactive surface hydroxyls. Recent publications on reactivity of O^- show the requisite high rate constants for abstraction of α^- hydrogen^{17, 25}.

Data to the left of the arrow in the diagrams in Fig. 7 were obtained prior to the commencement of u.v. illumination and show that presence of N_2O and isopropanol in 1:1 mole ratio as the gas phase flowing over the 'dark' ZnO interface resulted in a larger pre-illumination level of acetone formation (Fig. 7A upper plot). Data to the right of the arrow in the upper plot show that the initial photoenhanced formation of acetone was larger and was sustained for longer times in the presence of N_2O .

These observations are consistent with some production of additional O^- species from N_2O on the dark or illuminated ZnO interface and reaction of these O^- species to yield dehydrogenated products via reactions of type 3a and 3b. Data on the upper plot of Fig. 7A at times >60 min. illumination indicate a small nett increase in acetone, corresponding to some small rate of photoassisted dehydrogenation in the photostationary state via 3a and 3b. Nitrogen corresponding to this process was detected in the static photo-reactions (see Fig. 2A ii).

Data in Fig. 7B illustrate the photoassisted growth of both the dehydration product, propene, and the dehydrogenation product, acetone, above a uv-illuminated isopropanol/ TiO_2^* interface. Comparison of Figures 7A and 7B confirm the greater activity of the TiO_2 surface for these photoassisted elimination reactions involving isopropanol. Since ESR measurements on a previous study on the ZnO and TiO_2 samples used in this work showed much greater surface concentration of metal-excess centres on TiO_2 than on ZnO, such Ti^{3+} centres are suggested as the active sites responsible for the greater dehydration on TiO_2 .

II.2 *Comparison of photoassisted elimination reactions for primary, secondary and tertiary Butanols.*

summarises results of on-line mass analysis of photoproducts detected above uv-illuminated interfaces of TiO_2 with primary, secondary or tertiary butanol. Extent of observed photo-dehydration may be seen to lie qualitatively in the sequence expected on the basis of ease of alcohol dehydration, viz. tertiary $>$ secondary $>$ primary. Since a quantitative measure of the extent of each photoassisted elimination reaction was desirable, values

for the total amount of gaseous photo-products which desorbed from the illuminated interfaces over the observed time-intervals were derived using equation (5),

$$\text{Amount desorbed} = \frac{1}{ARTo} V\Delta P(t) + S \int_0^t \Delta p dt \quad (5)$$


In this equation, as developed in 1953 by Becker and Hartmann²⁶ A = surface area of adsorbent; To = gas temperature; V = volume of reaction system; S = pumping speed and $\int_0^t \Delta p dt$ is the area under pressure-versus-time curves such as are shown in figures 7 and 8. Increases in peak height obtained experimentally for various mass numbers were converted to pressure changes Δp , and substituted into equation (5), thereby yielding values for the total number of molecules of each product desorbing in a given time interval, (usually 0-170 min. after the commencement by uv. illumination). These values are tabulated in Table 4 for the Alcohol/Metal Oxide systems studied. In relation to photodehydration, these data show that extent of reaction of tertiary butanol was an order of magnitude greater than for secondary alcohols at the same interface, whilst primary alcohols did not experience photodehydration to measurable extent. Extent of photodehydrogenation (see columns headed "-H₂" in Table 4) did not show such large variations with the nature of the alcohols, which indicates that extent of this photoassisted elimination process depended more upon the number of active sites provided by the illuminated TiO₂ or ZnO surface than upon ease of alcohol dehydrogenation. Tertiary butanol represented an obvious exception to this behaviour since dehydrogenation via. an initial α -hydrogen abstraction was not possible. Our observations showed that acetone was the second major photoproduct from t-butanol/

TABLE 4

Extent of room temperature dehydration and dehydrogenation at alcohol vapour/
metal oxide illuminated interfaces under dynamic conditions.

Reactant	ZnO [*]			TiO ₂ [*]		
	(-H ₂ O)	(-H ₂)	(-CH ₄)	(-H ₂ O)	(-H ₂)	(-CH ₄)
C ₂ H ₅ OH	-	2.5x10 ¹⁹	-	-	1.1x10 ²⁰	-
N ₂ O+C ₂ H ₅ OH	-	6.9x10 ¹⁹	-	-	2.3x10 ²⁰	-
(CH ₃) ₂ CHOH	-	3.6x10 ¹⁹	-	1.9x10 ¹⁹	1.4x10 ²⁰	-
N ₂ O+(CH ₃) ₂ CHOH	-	7.7x10 ¹⁹	-	-	2.6x10 ²⁰	-
(CH ₃) ₃ COH	1.1x10 ²⁰	-	7.4x10 ¹⁹	1.8x10 ²⁰	-	1.8x10 ²⁰
N ₂ O+(CH ₃) ₃ COH	8.4x10 ¹⁹	-	9.9x10 ¹⁹	6.1x10 ¹⁹	-	3.4x10 ²⁰
(CH ₃) ₂ CHCH ₂ OH	-	2.9x10 ¹⁹	-	-	7.8x10 ¹⁹	-
CH ₃ CH ₂ CH(OH)CH ₃	1.0x10 ¹⁹	4.3x10 ¹⁹	-	1.5x10 ¹⁹	1.3x10 ²⁰	-

Metal Oxide* systems, in addition to isobutene. The overall stoichiometry of acetone formation may be expressed as in (6).



A reaction with this stoichiometry would be consistent with the relatively large amounts of methane obtained on illumination of the *t*-butanol/metal oxide interface in the static reaction system (see Table 3). Such photoassisted reaction may be described as "(-CH₄)", i.e. a photoelimination of methane. Data listed under that heading in Table 4 indicate the amount of acetone detected by on-line mass analysis. It was originally felt that the formation of acetone at the *t*-butanol/ZnO* interface might be the result of secondary reactions such as oxidation of isobutene photoproduct by lattice oxygen. However, test experiments with mixture of (N₂O+isobutene) present over illuminated ZnO or TiO₂ surfaces under either static or dynamic conditions did not result in significant photooxidation of the isobutene. Dehydration followed by oxidative cleavage of the double bond is therefore discounted as a route to photoassisted elimination of the methane observed from *t*-butanol. In effect the process has the characteristics of a reverse Grignard reaction, which suggests involvement of co-ordinatively unsaturated metal ion sites on the catalyst surface. Further studies will be necessary before the nature of the sites active for this process can be identified with confidence.

II.3 Influence of water vapour in dynamic conditions.

Since analysis with the on-line mass analyses offered a more sensitive procedure (than static techniques) for examining

the influence of coadsorbed water upon photocatalytic activity, comparison was made of the extent of photoassisted formation of CD_3CDO from anhydrous $\text{C}_2\text{D}_5\text{OD}$ and from $\text{C}_2\text{D}_5\text{OD}$ containing 5% D_2O . Results are summarised in Fig. 8 for both ZnO and TiO_2 surfaces exposed to continuous illumination at 254nm. It is clear from these data that the presence of D_2O enhanced the extent of alcohol photo dehydrogenation. This is similar to reports by Stone et al¹⁶ that rehydroxylation of a TiO_2 surface resulted in increased photodehydrogenation of isopropanol to acetone.

CONCLUSIONS

Results obtained in this study for the extent of N_2O dissociation at vacuum-activated ZnO or TiO_2 surfaces are fully consistent with the interpretation previously advanced that electron transfer to N_2O from unsaturated metal ion sites i.e., $\text{M}_{\text{cus}}^{+(z-1)} \xrightarrow{\text{N}_2\text{O}} (\text{M}^{+z} - \text{O}^-)_{\text{cus}} + \text{N}_2$, represents the main channel for nitrogen production at room temperature in the absence of uv-illumination. This model for surface reactivity is extended by the new data here presented on further dissociation of N_2O in the presence of primary or secondary alcohol, allied to the evidence for enhanced alcohol dehydrogenation when N_2O was present. Together, these observations are consistent with secondary reactions of the O^- ions formed from N_2O in this process. A necessary corollary to its efficient occurrence in the presence of N_2O is that other reactions requiring $\text{M}_{\text{cus}}^{+(z-1)}$ sites should be inhibited. The inhibition of alcohol

dehydration here observed at surfaces also exposed to N_2O may thus be understood.

Results obtained here on photoassisted elimination reactions from alcohol vapours at ZnO or TiO_2 surfaces exposed to uv-illumination in static or dynamic conditions demonstrate the occurrence of other surface sites capable of interacting directly with alcohol when activated by illumination. Results with alcohols are consistent with identification of these sites as O_{cus}^{2-} and with their conversion to O_{cus}^- by hole-trapping during illumination. The enhancing effect of 5% D_2O upon photo-dehydrogenation of C_2D_5OD in dynamic conditions would be consistent with D_2O acting as an electron trap to complement such hole trapping. Present results do not, however, support the proposal that adsorbed alcohol act as efficient hole traps on ZnO or TiO_2 , even when N_2O was simultaneously present as an electron trap.

Results here presented on changes in surface reactivity effected by treatments designed to rehydroxylate the vacuum-activated ZnO or TiO_2 surfaces point to important differences between the sites so produced on TiO_2 from those on ZnO , with only the latter effective for N_2O dissociation. Further study will be necessary to fully characterise such sites and determine if they are in agreement with recent suggestions²⁷ that O_{cus}^{2-} adjacent to surface OH^- can act as electron donor sites on ZnO and NO_2 .

References

1. Work supported in part by funds from the U.S. Air Force Office of Scientific Research through EOAR under grants nos. 71-2148 and 76-2986.
2. Cunningham, J., Kelly, J.J., and Penny, A.L., *J. Phys. Chem.*, 74, (1970), 1992.
3. Cunningham, J., Kelly, J.J., and Penny, A.L., *J. Phys. Chem.*, 75, (1971), 617.
4. Cunningham, J., and Penny, A.L., *J. Phys. Chem.*, 76, (1972), 2353.
5. Cunningham, J., and Penny, A.L., *J. Phys. Chem.*, 78, (1974), 870.
6. Dewald, J.F. (a) *Bell System Tech. J.*, 39, (1960), 615 (b) *J. Phys. Chem. Solid*, 14, (1960), 155. (c) Cunningham, J. and Corkery, S., *J. Phys. Chem.*, 79, (1975), 933.
7. Lundsford, J.H., Taarit, Y.B. and Wong, N.B., *J. Chem. Phys.*, 60, (1974), 2148.
8. Che, M., Naccache, C. and Imelik, B., *J. Catalysis*, 24, (1972), 328.
9. Hosaka, H., Fujiwara, T. and Meguro, K., *Bull Chem. Soc. Japan*, 44, (1971), 2616.
10. Pimenov, Y.D., Kholmogorov, V.E., and Terenin, A.N., *Doklady Akad Nauk S.S.S.R.*, 163, (1965), 935.
11. (a) Ueda, H. *Bull Chem. Soc. Japan*, 43, (1970), 319
(b) Kiseler, A.V. and Urarov, A.V., *Surface Science*, 6, (1967), 399.
12. Davison, R.S. and Slater, R.M., *J.C.S. Farad. Trans.*, I, 72, (1976), 2416.
13. Morrison, S.R. and Freund, T., *Electrochem Acta.*, 13, (1968), 1343.
14. Gomes, W.P., Freund T. and Morrison, S.R., *J. Electrochem. Soc.*, 115, (1968), 818.
15. Bickeley, R.I. and Janaty, R.K.M., *Disc Farad Soc.*, 58, (1974), 194.
16. Bickeley, R.I. and Stone, F.S., *J. Catal*, 31, (1973), 389, 398.
17. Warman, J.M., *J. Phys. Chem.*, 72, (1968), 52.
18. Warman, J.M., *Nature*, 213, (1967), 381.
19. Atherton, K., Hockey, J.A. and Newbold Co., *Disc Farad. Soc.*, 52, (1971), 33.
20. Tench, A. and Nelson, R.L., *Trans. Faraday Soc.*, 63, (1967), 2254.
21. Primet, M., Pichat, P. and Mathieu, M.C., *J. Phys. Chem.*, 75, (1971), 1216.
22. Wolkenstein, T.L., *Electron Theory of Catalysis on Semiconductors* (Pergamon Press 1963).
23. Hauffe, K., *Adv. Catal*, 7, (1955), 213.
24. McArthur, D., Bliss, H. and Butt, J., *J. Catal*, 28, (1973), 183.
25. Zehani, D. and Rabani, J., *J. Phys. Chem.*, 75, (1971), 1738.
26. Becker, J.A. and Hartmann, C.D., *J. Phys. Chem.*, 57, (1953), 157.
27. Cordischi, D. and Indovina, V., *J.C.S. Farad. Trans.*, I, 72, (1976), 2341.

Captions to Figures

Figure 1

Vacuum systems for study of gaseous products evolved from interfaces of ZnO and TiO₂ with N₂O and/or aliphatic alcohols:

1(a) Static reactor with pumping line, P; gas handling line, L; pirani gauge, G; variable leak valve, V; metal valves, M; cold fingers, C; and metal oxide sample, M.O.

1(b) Dynamic reactor with inlet leak valve, I; metal oxide layer, M.O.; stainless steel tubing, E; glass walled photo-reactor, C; micromass 6 mass analyser, M.A.; pressure gauge, B; and ion pump.

Figure 2

Evolution of N₂ product from ZnO surfaces at 293K (position of arrow denotes commencement of illumination):

- 2A (i) N₂O/ZnO with $P_{N_2O} = 45 \text{ Nm}^{-2}$.
(ii) N₂O + C₂H₅OH/ZnO with $P_{N_2O} = 45 \text{ Nm}^{-2}$
 $P_{C_2H_5OH} = 45 \text{ Nm}^{-2}$.
- 2B (i) N₂O + (CH₃)₃COH/ZnO with $P_{N_2O} = 45 \text{ Nm}^{-2}$
 $P_{(CH_3)_3COH} = 45 \text{ Nm}^{-2}$.
(ii) N₂O + (CH₃)₂CHOH/ZnO with $P_{N_2O} = 45 \text{ Nm}^{-2}$
 $(CH_3)_2CHOH = 45 \text{ Nm}^{-2}$.

Figure 3

Effects of surface rehydroxylation upon N₂O dissociation on ZnO or TiO₂ at 293K:

- 3A (i) N₂ evolution on ZnO surface pretreated in vacuo at 623K for 16 hours (arrow denotes start of illumination).
(ii) As for (i) except surface rehydroxylated by treatment in 50 Nm⁻² H₂O at 623K for 2 hours.
(iii) As for (ii) except u.v. illumination commenced at time indicated by the arrow.
- 3B (i) N₂ from TiO₂ surface pretreated in vacuo at 623K for 16 hours
(ii) As for (i) except surface rehydroxylated at 623K in 0.5 torr H₂O.

Figure 4

Evolution of N_2 product from TiO_2 surfaces at 293K:

- 4A (i) N_2O/TiO_2 ;
(ii) $N_2O + C_2H_5OH/TiO_2$;
- 4B (i) $N_2O + (CH_3)_3COH/TiO_2$;
(ii) $N_2O + (CH_3)_2CHOH/TiO_2$.

Figure 5

Histograms for thermal desorption of various products from $(CH_3)_2CHOH/ZnO$ interfaces on heating to 623K:

- (i) H_2 evolution; (ii) propene;
(iii) Acetone (darkened sections represents thermal desorption of $(CH_3)_2CO$ from a $(CH_3)_2CO/ZnO$ interface.

Figure 6

E.S.R. spectra of TiO_2 at 77K:

- (i) After activation for 16 hours at 77K.
(ii) After exposure for 10 min. to C_2H_5OH at 293K followed by evacuation.

Figure 7



Time profiles for appearance of various products in the gas-phase, as detected with Micromass 6 mass analyser with isopropanol flowing over the metal-oxide catalyst at $2 \times 10^{-4} Nm^{-2}$. Arrows denote the start of continuous illumination with photons at 254 n.m.


- 7A (i) Acetone from isopropanol/ ZnO^* .
(ii) Acetone from $N_2O +$ isopropanol/ ZnO^* .
- 7B (i) Acetone from isopropanol/ TiO_2^*



Figure 8

Influence of a 5% water content upon photodehydrogenation of ethanol over ZnO and TiO_2 surfaces illuminated by 254 n.m. photons in a dynamic quartz photoreactor.


- 8A (i) CD_3CDO from C_2D_5OD/ZnO^*
(ii) CD_3CDO from $C_2D_5OD + 5\%D_2O/ZnO^*$.
- 8B (i) CD_3CDO from C_2D_5OD/TiO_2^*
(ii) CD_3CDO from $C_2D_5OD + 5\%D_2O/TiO_2^*$.













$V_n \times 10^{-4} / \text{ml s per metre}^2$











ATE
LME